
2.8.2023

Cyanobacterial hepatotoxin(Microcystin) concentration in Littoistenjärvi 2023







On 2.8.2023, microcystin concentration in Littoistenjärvi water is very low (below detection limit of 0.25 μ g/L).

Interpretation (3.8.2023 SA)

Collection of Raw water samples: 2.8.2023,

Immunoassay analysis: 3.8.2023

Immunoassay detection limit on 3.8.2023: 0.25 μg/L of Microcystin-LR

Before analysis, samples were heated at 90 °C for 10 min to release cell bound toxins if any.

The results represent the total cyclic peptide hepatotoxin amount (already released toxin in water and the cell bound toxin) .

The immunoassay detects cyanobacterial peptide hepatotoxins (microcystins and/or nodularin).

For quantification, microcystin-LR was used as standard.

Result:

In Littoistenjärvi water, the detected cyanobacterial peptide hepatotoxin (free and cell bound microcystin) concentrations (μ g/L) are as follows:

2.8.2023 A Saarten taus: below 0.25 μg/L

B_Koilliselkä: below 0.25 μg/L C_Luoteisselkä: below 0.25 μg/L A' Hiekkaranta: below 0.25 μg/L

D'_Ristikallion Uimaranta: below 0.25 μg/L

World Health Organization (WHO) recommended provisional guideline values

Provisional lifetime **drinking**-water guideline value for microcystin: $1 \mu g/L$

Provisional short-term (2 weeks) drinking-water guideline value for microcystin 12 μ g/L (Adult), 3 μ g/L (children)

Provisional **recreational** water guideline value for microcystin: **24 μg/L**

Reference:

Chorus, I., & Welker, M. (2021). Toxic cyanobacteria in water: a guide to their public health consequences, monitoring and management (p. 858). Taylor & Francis. P25-28

Date of analysis: 3.8.2023
Report by Sultana Akter,
Project Researcher,
Department of Life Technologies (Biotechnology),
Faculty of Technology, University of Turku