

# 20.7.2023

Cyanobacterial hepatotoxin(Microcystin) concentration in Littoistenjärvi 2023



On 20.7.2023, microcystin concentration in Littoistenjärvi water is very low (below 0.2 μg/L)





## Interpretation (21.7.2023 SA)

Collection of Raw water samples: 20.7.2023,

Immunoassay analysis: 21.7.2023

Before analysis, samples were heated at 90 °C for 10 min to release cell bound toxins if any.

The results represent the total cyclic peptide hepatotoxin amount (already released toxin in water and the cell bound toxin) .

The immunoassay detects cyanobacterial peptide hepatotoxins (microcystins and/or nodularin).

For quantification, microcystin-LR was used as standard.

#### **Result:**

In Littoistenjärvi water, the detected cyanobacterial peptide hepatotoxin (free and cell bound microcystin) concentrations ( $\mu$ g/L) are as follows:

**20.7.2023** A\_Saarten taus: below 0.2 μg/L

B\_Koilliselkä: below 0.2 μg/L C\_Luoteisselkä: below 0.2 μg/L A' Hiekkaranta: below 0.2 μg/L

D' Ristikallion Uimaranta: below 0.2 µg/L

# World Health Organization (WHO) recommended provisional guideline values

Provisional lifetime **drinking**-water guideline value for microcystin:  $1 \mu g/L$ 

Provisional short-term (2 weeks) drinking-water guideline value for microcystin 12  $\mu$ g/L (Adult), 3  $\mu$ g/L (children)

Provisional **recreational** water guideline value for microcystin: **24 µg/L** 

### Reference:

Chorus, I., & Welker, M. (2021). Toxic cyanobacteria in water: a guide to their public health consequences, monitoring and management (p. 858). Taylor & Francis. P25-28

Date of analysis: 21.7.2023

Report by Sultana Akter, Project Researcher, Department of Life Technologies (Biotechnology), Faculty of Technology, University of Turku